

# High $p_T$ Spectra of Identified Particles Produced in Pb+Pb Collisions at 158 GeV/nucleon Beam Energy

András László<sup>a</sup> and Tim Schuster<sup>b</sup> for the  Collaboration\*

<sup>a</sup>MTA-KFKI Research Institute of Particle and Nuclear Physics, Budapest, Hungary

<sup>b</sup>Institute for Nuclear Physics, J. W. Goethe University, Frankfurt am Main, Germany

Transverse momentum spectra of  $\pi^\pm$ ,  $p$ ,  $\bar{p}$ ,  $K^\pm$ ,  $K_s^0$  and  $\Lambda$  at midrapidity were measured at high  $p_T$  in Pb+Pb collisions at 158 GeV/nucleon beam energy by the NA49 experiment. Particle yield ratios ( $p/\pi$ ,  $K/\pi$  and  $\Lambda/K_s^0$ ) show an enhancement of the baryon/meson ratio for  $p_T > 2$  GeV/c. The nuclear modification factor  $R_{CP}$  is extracted and compared to RHIC measurements and pQCD calculations.

## 1. INTRODUCTION

One of the most interesting features discovered at RHIC is the suppression of high  $p_T$  particle production in central nucleus-nucleus reactions relative to peripheral ones or p+p collisions. This is generally interpreted as a sign of parton energy loss in hot and dense nuclear matter. Additionally, an enhancement of baryon/meson ratios above unity at high  $p_T$  was observed and can be explained in the context of quark coalescence models. The aim of this analysis is to investigate the energy dependence of these effects by studying nucleus-nucleus reactions at top SPS energy ( $\sqrt{s_{NN}} = 17.3$  GeV/nucleon) (see: [1,8]).

## 2. DATA ANALYSIS

Centrality selection is based on a calorimetric measurement of the energy observed in the projectile spectator region of phase space (see [3]). Charged particle spectra ( $\pi^\pm$ ,  $p$ ,  $\bar{p}$  and  $K^\pm$ ) in the center of mass rapidity interval  $[-0.3, 0.7]$  are analyzed in the centrality ranges (0-5)%, (12.5-23.5)%, (33.5-80)% of the total inelastic cross section. The tracking efficiency for single tracks is above 95% and an efficient fake track rejection is applied. The particle identification is done by unfolding the energy loss spectra measured in different phase-space bins. The typical  $\frac{dE}{dx}$  resolution varies between 3 and 6%. The  $\pi^\pm$  and  $p$ ,  $\bar{p}$  yields were not corrected for feed down from the decay of  $K_s^0$  and hyperons; furthermore the  $K^\pm$  yields were not corrected for decay loss.

Neutral strange particles were analyzed in the centrality range (0-23.5)%. They are identified via the topology of their weak decay into the channels  $K_s^0 \rightarrow \pi^+\pi^-$  (BR = 68.95%) and  $\Lambda \rightarrow p\pi^-$  (63.9%). For the V0-candidates, selected by geometrical criteria,

---

\*For the full list of NA49 authors and acknowledgments, see appendix 'Collaborations' of this volume.

the invariant mass of the daughter particles is calculated as a function of  $p_T$  and the yields of  $K_s^0$  and  $\Lambda$  are extracted on a statistical basis. The shown results are for the rapidity interval  $[-0.5, 0.5]$  and corrected for acceptance and reconstruction inefficiency. The  $\Lambda$  yields are not corrected for feed down from the decay of heavier hyperons.

### 3. PHYSICS RESULTS

The proton/pion and the kaon/pion ratios are shown in Fig. 1. These ratios exhibit a monotonic increase with  $p_T$  and centrality at high  $p_T$ . The kaon/pion ratios show a saturation tendency at high  $p_T$ , particularly the  $K^-/\pi^-$  ratio.




Figure 1. Proton/pion (upper panels) and kaon/pion (lower panels) ratios vs.  $p_T$  and centrality.

In the left panel of Fig. 2, our measurement of proton/pion ratio is compared to RHIC data. The shape of these curves is approximately energy independent. The right panel of Fig. 2 shows NA49 baryon/meson ratios, compared to a Blast-Wave (BW, see [7]) parametrization of  $m_T$  spectra and radius parameters from Bose-Einstein correlations of pions, fitted simultaneously at low  $p_T$ . The BW model curve does not describe the data at high  $p_T$ .

The nuclear modification factor  $R_{CP}$  is defined by  $R_{CP} := \frac{N(\text{Peripheral})}{N(\text{Central})} \cdot \frac{\text{Yield}(\text{Central})}{\text{Yield}(\text{Peripheral})}$ . Here  $N$  can be either the number of binary collisions or the number of wounded nucleons obtained from model calculations in the given centrality range. The upper panels of Fig.




Figure 2. The energy dependence of proton/pion ratios (left panel), and a comparison of the baryon/meson ratios at top SPS energy to a Blast-Wave parametrization (right panel).

3 show the energy dependence of  $R_{CP}$  vs.  $p_T$  of pions with binary collision and with wounded nucleon scaling. At high  $p_T$  there is a strong energy dependence with both scalings, however at low  $p_T$  wounded nucleon scaling makes  $R_{CP}$  energy independent. A similar phenomenon was pointed out for unidentified particles in [6]. The lower panels of Fig. 3 show the comparison of our data to pQCD calculations (see [10]).  $R_{CP}$  is consistent with the pQCD calculation at  $p_T > 2$  GeV/c. However, the pQCD prediction for the antibaryon/meson ratio is very far from the data below 4 GeV/c.

#### 4. CONCLUDING REMARKS

First NA49 results on particle yields around midrapidity in the range  $2 \text{ GeV}/c \leq p_T < 4.5 \text{ GeV}/c$  were presented from a study of 158 GeV/nucleon beam energy Pb+Pb collisions.

A monotonic increase of baryon/meson ratios and kaon/pion ratios with  $p_T$  and centrality was observed at high  $p_T$ . The  $p_T$  shape of the baryon/meson ratio is approximately energy independent. The measured baryon/meson ratios were compared to a Blast-Wave model: the model predictions exceed the data for  $p_T > 1.5$  GeV/c.

The nuclear modification factors  $R_{CP}$  were also determined from the particle yields for various particle species, as a function of  $p_T$ . The measured  $R_{CP}$  ratio does not show Cronin enhancement for the mesons at larger  $p_T$  when using binary collision scaling. The behavior is qualitatively similar to the  $p_T$  shape observed at RHIC. A strong energy dependence of the  $R_{CP}$  ratios was observed at high  $p_T$  with both binary collision and wounded nucleon scaling. However, at low  $p_T$ , the wounded nucleon scaling factorizes out the energy dependence. Results for  $R_{CP}$  with binary collision scaling are consistent with pQCD model calculations at  $p_T > 2.5$  GeV/c. However, the pQCD calculation strongly overpredicts the observed antibaryon/meson ratio for  $p_T < 4$  GeV/c.




Figure 3. Energy dependence of  $R_{CP}$  vs.  $p_T$  (upper panels): binary collision scaling (left panel) and wounded nucleon scaling (right panel). Comparison of data to pQCD calculations (lower panels): the nuclear modification factor  $R_{CP}$  (left panel) and the baryon/meson ratio (right panel).

## REFERENCES

1. S. S. Adler *et al* (the PHENIX Collaboration): Phys. Rev. C69, 034909, 2004.
2. S. S. Adler *et al* (the PHENIX Collaboration): Phys. Rev. C69, 034910, 2004.
3. A. Afanasiev *et al* (the NA49 collaboration): Nucl. Instr. Meth., A430, 210-244, 1999.
4. A. Afanasiev *et al* (the NA49 collaboration): Phys. Rev., C66, 054902, 2002.
5. T. Anticic *et al* (the NA49 collaboration): Phys. Rev., C69, 024902, 2004.
6. B. B. Back *et al* (the PHOBOS Collaboration): Phys. Rev. Lett., 94, 082304, 2005.
7. F. Retiere, M. A. Lisa: Phys. Rev., C70, 044907, 2004.
8. L. Ruan (for the STAR Collaboration): J. Phys., G31, S1029-S1033, 2005.
9. M. Shao (for the STAR Collaboration): J. Phys., G31, S85-S92, 2005.
10. X.-N. Wang: Phys. Lett., B595, 165-170, 2004.